

Reply to Comment on “Surprises in threshold antikaon-nucleon physics”

José A. Oller¹, Joaquim Prades² and Michela Verbeni¹

¹ *Departamento de Física, Universidad de Murcia, E-30071 Murcia, Spain*

² *Centro Andaluz de Física de las Partículas Elementales (CAFPE) and Departamento de Física Teórica y del Cosmos, Universidad de Granada, Campus de Fuente Nueva, E-18002 Granada, Spain*

In their Comment, Borasoy *et al.* [1], criticize our results [2] that accommodate both scattering data and the new accurate measurement by DEAR of the shift and width of kaonic hydrogen. In our calculations we have employed unitary chiral perturbation theory (UCHPT). We discuss why their arguments are irrelevant or do not hold.

1. Borasoy *et al.* correctly state in their Comment that the interacting kernel employed in the Letter [2] can induce residual cuts in some energy regions above the threshold of the lighter channels ($\pi\Lambda$ and $\pi\Sigma$) because of u -crossed exchange of baryons in amplitudes with heavier states. However, these cuts only induce appreciable unitarity deviations in $\pi^0\Lambda \rightarrow \pi^0\Lambda$ at energies lower than 1.4 GeV, where no data exists to compare with. These deviations are negligible for the $\pi\Sigma$ and do not appear at all in the rest of channels. UCHPT resums the right-hand cut (or unitarity cut) while the contributions from the other cuts in the interacting kernel are calculated only at a given order in CHPT, by matching with the chiral series order by order. The right hand cut resummation is justified since the chain of unitarity bubbles is enhanced by the large masses of kaons and baryons [3]. In this way, one reproduces the CHPT results up to a given order, e.g. to $\mathcal{O}(p^2)$ in [2]. In this scheme, calculating the interacting kernel at higher orders in CHPT will eventually soften the effect of these spurious cuts. If one eliminates the u -channel cuts in the S-wave projections by ad hoc taking them to be constants above some energy value, as in [4], then *chiral symmetry as well as crossing are violated* even at leading order in CHPT.

2. Borasoy *et al.* [1] consider necessarily unphysical the narrow pole just below the K^-p threshold at (1431.21- i 1.28) MeV. First of all, one must point out that this pole does not occur in the *physical* Riemann sheet but in the unphysical ones as required by hermiticity. Furthermore, the fact that this resonance turns out to be so narrow has a clear physical origin and is not an artifact at all. This resonance has couplings (with respect to $I = 1$ states) of (1.8,4.5,1.0,0.7,0.5) GeV to the ($K\Xi, \eta\Sigma, \bar{K}N, \pi\Sigma, \pi\Lambda$), respectively. One could be surprised that the couplings to heavier channels (the $\eta\Sigma$ threshold is around 200 MeV above the resonance mass) are larger than those to the channels closer in energy. But this is also the case for the $\Lambda(1670)$, well known experimentally, that couples much more strongly to the $K\Xi$ channel [5] than to the rest of channels. The pole at 1431 MeV is so narrow because of the smallness of its couplings to the open channels. They also argue that such narrow pole should lead to violations of fundamental principles like the Wigner condition. These authors

apply such condition to the phase shifts. However in the inelastic case (as the one here), one should consider the phase of the scattering amplitude. This increases the derivative from -300 [1] to -150 fm and makes Figure 1 in [1] irrelevant to the present discussion since inelasticity varies sharply. Furthermore, the Wigner condition is not violated in the presence of a narrow pole, as the derivative of the phase is positive (take a Breit-Wigner) and the Wigner condition is a lower negative bound. The phase of the scattering amplitude starts to decrease with energy between the pole and the opening of the K^-p channel at 1431.95 MeV due to the presence of the branch singularity associated with the opening of this channel. The Wigner condition cannot be applied on top of the thresholds of new opening channels, simply because the phase is not differentiable in an inelastic branch point and the derivative near it tends to infinity. If the situation is softened, by varying the masses a few MeV so that the K^-p threshold moves away, then the Wigner limit is restored.

3. The authors of [1] claim that this pole should be seen in the $\pi\Sigma$ event distribution. However, this is not so because the $\pi\Sigma$ distribution is dominated by $I = 0$ and this narrow resonance is $I = 1$. But even if we use the $K^-p \rightarrow \pi^\pm\Sigma^\mp$ amplitudes one does not observe any narrow peak because of the interference with $I = 0$, which is larger than $I = 1$, as we have checked. One would need much better precision than in present data to observe the modifications in the event distribution due to such pole. Finally, the authors of [1] also state that our fit is too sensitive to the values of the fitted parameters. One must remark that the set of parameters is highly correlated so it does not make any sense to arbitrarily vary their values. What is sensible is to change the set of fitted data, e.g. by modifying the values of $\sigma_{\pi N}$, m_0 or by considering new data points, and check that similar fits arise.

-
- [1] B. Borasoy, R. Nißler and W. Weise, hep-ph/0512279.
[2] J.A. Oller, J. Prades and M. Verbeni, Phys. Rev. Lett. **95**, 172502 (2005).
[3] S. Weinberg, Nucl. Phys. **B363**, 3 (1991).
[4] B. Borasoy, R. Nißler and W. Weise, Phys. Rev. Lett. **94**,

- 213401 (2005); Eur. Phys. J. **A25**, 79 (2005).
[5] D. Jido *et al.*, Nucl. Phys. **A725**, 181 (2003).